

ARISTADA
INITIO®
aripiprazole lauroxil
extended-release injectable suspension

675 mg

ARISTADA®
aripiprazole lauroxil
extended-release injectable suspension

441 mg

662 mg

882 mg

1064 mg

Images are not
actual patients


FOR CAREGIVERS OF
ADULTS WITH SCHIZOPHRENIA

Find out if a treatment option given

EVERY 2 MONTHS*
(1064 mg)

may be right for your loved one

Other dosing options available



*After establishing tolerability to oral aripiprazole and completing starting regimen.

Scan this QR code using your smartphone camera to watch a video about ARISTADA



INDICATION

ARISTADA INITIO® (aripiprazole lauroxil) is a prescription medicine given as a **one-time injection** and is used in combination with oral aripiprazole to start ARISTADA® (aripiprazole lauroxil) treatment, or re-start ARISTADA treatment after a missed dose, when ARISTADA is used for the treatment of schizophrenia in adults.

ARISTADA is a prescription medicine given by injection by a healthcare professional and used to treat schizophrenia in adults.

It is not known if ARISTADA INITIO and/or ARISTADA are safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA

Elderly people with dementia-related psychosis are at increased risk of death when treated with antipsychotic medicines including ARISTADA INITIO and ARISTADA. ARISTADA INITIO and ARISTADA are not for the treatment of people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia).

Please see additional Important Safety Information throughout and full Prescribing Information for [ARISTADA INITIO](#) and [ARISTADA](#), including Boxed Warning and Medication Guides for [ARISTADA INITIO](#) and [ARISTADA](#).

KNOW SOME SIGNS of Schizophrenia

Certain behaviors could be signs that someone may have schizophrenia. These include:

- Social withdrawal
- Hostility or suspiciousness
- Lack of expression
- Decline in personal hygiene
- Trouble focusing or paying attention

These are not all of the signs of schizophrenia

It's important to talk to a healthcare provider if you are worried that someone you love may have schizophrenia.



Please see additional Important Safety Information throughout and full Prescribing Information for **ARISTADA INITIO** and **ARISTADA**, including Boxed Warning and Medication Guides for **ARISTADA INITIO** and **ARISTADA**.

Is a LONG-ACTING TREATMENT right for your loved one?

The most common medicines for schizophrenia can be taken by mouth or by injection. Both of these options provide the treatment needed. The difference is how they're taken.



LONG-ACTING INJECTABLES

are given by a healthcare provider and can offer different durations of treatment, depending on the medication chosen.



ORAL MEDICATIONS,

such as pills, are taken by mouth every day.

Help the person you care for explore their treatment options

There are multiple options for treating schizophrenia. It can be helpful to find out what treatment options are available. Then, you and/or your loved one can have an informed discussion with their healthcare provider about which treatment approach may be right for them.



IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (*continued*)

Contraindication: Do not receive ARISTADA INITIO or ARISTADA if you are allergic to aripiprazole or any of the ingredients in ARISTADA. Allergic reactions to aripiprazole have ranged from rash, hives and itching to anaphylaxis, which may include difficulty breathing, tightness in the chest, and swelling of the mouth, face, lips, or tongue.

ARISTADA INITIO and/or ARISTADA may cause serious side effects including:

- **Cerebrovascular problems (including stroke) in elderly people with dementia-related psychosis that can lead to death.**

1 dose can provide treatment for 2 MONTHS

ARISTADA® is a long-acting injectable medication that can provide 2 months (1064 mg) of treatment. When receiving ARISTADA, your loved one will have one less medicine to remember to take every day.

After establishing tolerability to oral aripiprazole, treatment with ARISTADA can start in one of two ways:

1. They will receive one injection of ARISTADA INITIO® (aripiprazole lauroxil) and one 30 mg dose of oral aripiprazole on the same day. The first injection of ARISTADA will be given on that same day or within the following 10 days.

OR

2. On the first day, they will receive an injection of ARISTADA and then take oral aripiprazole daily for 21 days if ARISTADA INITIO is not right for them.

It's important that they not miss a dose of ARISTADA. If a dose is missed, they should contact their healthcare provider to discuss what to do next.

See if a 2-MONTH dosing option may be right for your loved one

How often they are treated with ARISTADA (aripiprazole lauroxil) depends on the dosage of medication that they and their healthcare provider decide is right for them.

Other dosing options include:

Every 2 months	Every 6 weeks	Every 1 month
1064 mg	882 mg	441 mg 662 mg 882 mg

ARISTADA is slowly released into the body to last for weeks at a time with 1 injection.

Please see additional Important Safety Information throughout and full Prescribing Information for ARISTADA INITIO and ARISTADA, including Boxed Warning and Medication Guides for ARISTADA INITIO and ARISTADA.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)


ARISTADA INITIO and/or ARISTADA may cause serious side effects including: (continued)

- **Neuroleptic malignant syndrome (NMS), a serious condition that can lead to death.** Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS:
 - high fever
 - sweating
 - stiff muscles
 - changes in pulse, heart rate, and blood pressure
 - confusion
- **Uncontrolled body movements (tardive dyskinesia).** ARISTADA INITIO and ARISTADA may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop receiving ARISTADA. Tardive dyskinesia may also start after you stop receiving ARISTADA.

What 1064 mg of ARISTADA (aripiprazole lauroxil)

EVERY 2 MONTHS

can look like:



6x only per year

WHAT TO KNOW

before your loved one starts treatment

- Before starting treatment with ARISTADA INITIO® (aripiprazole lauroxil) or ARISTADA® (aripiprazole lauroxil), tell their healthcare provider if they have not taken Abilify®, Abilify Maintena®, or any aripiprazole product before
 - This is because people who are allergic to aripiprazole or any of the ingredients in ARISTADA should not receive ARISTADA or ARISTADA INITIO
- If they've never taken aripiprazole before, their healthcare provider may have them take oral aripiprazole for a short time to see if they can tolerate it before they start treatment

It's also important to tell the healthcare provider about all their medical conditions, including if they:

- Have or had heart problems or stroke
- Have diabetes or high blood sugar or a family history of diabetes or high blood sugar. The healthcare provider should check on their blood sugar before they receive ARISTADA INITIO and/or ARISTADA and during their treatment
- Have or had low or high blood pressure
- Have or had seizures (convulsions)
- Have or had a low white blood cell count
- Have problems that may affect them receiving an injection in their buttocks or their arm
- Are pregnant or plan to become pregnant. It is not known if ARISTADA INITIO and/or ARISTADA will harm their unborn baby. If they become pregnant while on treatment, talk to the healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. They can register by calling 1-866-961-2388, or visit womensmentalhealth.org/research/pregnancyregistry/
- Are breastfeeding or plan to breastfeed. ARISTADA INITIO and/or ARISTADA can pass into their breast milk. It is not known if it may harm their baby. Talk to the healthcare provider about the best way to feed their baby if they receive ARISTADA INITIO and/or ARISTADA.

Please see additional Important Safety Information throughout and full Prescribing Information for **ARISTADA INITIO** and **ARISTADA**, including **Boxed Warning** and **Medication Guides** for **ARISTADA INITIO** and **ARISTADA**.

Demonstrated in clinical studies to reduce SYMPTOMS OF SCHIZOPHRENIA in adult patients

Every person responds to schizophrenia medication differently. In clinical studies, ARISTADA® (aripiprazole lauroxil) was shown to reduce symptoms of schizophrenia.

Some common symptoms:

- Hallucinations
- Delusions
- Unusual thinking or disorganized speech
- Loss of motivation
- Disinterest or lack of enjoyment in daily activities
- Difficulty showing emotions
- Difficulty processing information to make decisions

These are not all of the symptoms of schizophrenia.

ARISTADA was shown to reduce symptoms of schizophrenia.

Ask their healthcare provider if ARISTADA

EVERY 2 MONTHS
(1064 mg)

could be right for your loved one.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)

ARISTADA INITIO and/or ARISTADA may cause serious side effects including: (continued)

- **Problems with your metabolism such as:**
 - **High blood sugar (hyperglycemia).** Increases in blood sugar can happen in some people who receive ARISTADA INITIO and/or ARISTADA. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes), your healthcare provider should check your blood sugar before you receive ARISTADA INITIO and/or ARISTADA and during your treatment with ARISTADA.



SUPPORT THEM
IN TAKING IT
ONE STEP
AT A TIME



Hear from real people
who received ARISTADA at
aristada.com/real-stories

Please see additional Important Safety Information throughout and full Prescribing Information for **ARISTADA INITIO** and **ARISTADA**, including Boxed Warning and Medication Guides for **ARISTADA INITIO** and **ARISTADA**.

GETTING STARTED

can happen in just one day

After discussing with their healthcare provider and establishing tolerability to oral aripiprazole, there are **3 steps to beginning their treatment:**

- 1.** The healthcare provider will give them a one-time injection of ARISTADA INITIO® (aripiprazole lauroxil) and a single 30 mg dose of oral aripiprazole.

Alternatively, their healthcare provider could give them oral aripiprazole for 21 days in a row when they receive their first injection of ARISTADA® (aripiprazole lauroxil).
- 2.** They will receive their first dose of ARISTADA either on the same day as ARISTADA INITIO or up to 10 days later.
- 3.** After receiving their initial doses, their following injections of ARISTADA (1064 mg) will be every 2 months, unless they and the healthcare provider have decided on a different dosing option.

How ARISTADA is given:

- Both ARISTADA INITIO and ARISTADA are given by injection by a healthcare provider into the muscle (intramuscular)
- Depending on the dose, ARISTADA can be given in the upper arm (441 mg only) or buttock (all doses). ARISTADA INITIO can be given in either the upper arm or buttock
- Follow the ARISTADA treatment schedule exactly as the healthcare provider says to
- There are multiple dosing options for ARISTADA. Talk to the healthcare provider to see what's right for them

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (*continued*)

ARISTADA INITIO and/or ARISTADA may cause serious side effects including: (*continued*)

- **Call your healthcare provider if you have any of these symptoms of high blood sugar:**
 - feel very thirsty
 - need to urinate more than usual
 - feel very hungry
 - feel weak or tired
 - feel sick to your stomach
 - feel confused, or your breath smells fruity
- **Increased fat levels (cholesterol and triglycerides) in your blood**
- **Weight gain.** You and your healthcare provider should check your weight regularly.

Safety has been evaluated
IN 1,180 PATIENTS*

ARISTADA® may cause side effects

The most common side effect of ARISTADA is restlessness, also known as akathisia.

Other side effects are possible, some of them serious. Please see additional Important Safety Information throughout this brochure. You can also review the Medication Guides for ARISTADA and ARISTADA INITIO® for more information.

Remember to talk to a healthcare provider about any questions or concerns you or your loved one have about starting treatment with **ARISTADA**.



Find a healthcare provider
in your area who can administer
ARISTADA INITIO (aripiprazole lauroxil) and
ARISTADA (aripiprazole lauroxil) at:



aristada.com/find-provider

*In clinical studies, 1,180 patients treated with ARISTADA were studied.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)

ARISTADA INITIO and/or ARISTADA may cause serious side effects including: (continued)

- **Unusual and uncontrollable (compulsive) urges.** Some people taking aripiprazole have had strong unusual urges such as gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having unusual strong urges, talk to your healthcare provider.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.

ARISTADA Care Support is
HERE TO HELP YOU
and your loved one



- If they have commercial insurance, they may be able to lower their out-of-pocket cost of treatment with ARISTADA® (aripiprazole lauroxil) through the ARISTADA Co-pay Savings Program*
- Patient Assistance Program for those who are eligible*
*Eligibility requirements and terms and restrictions apply
- Help finding a healthcare provider who can administer ARISTADA INITIO® (aripiprazole lauroxil) and/or ARISTADA through the ARISTADA Provider Network

To speak to an ARISTADA
Care Support representative, call
1-866-ARISTADA (1-866-274-7823),
Monday through Friday, 9 AM to 8 PM ET.

Visit aristada.com/resources for more helpful links and information about living with schizophrenia

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)

ARISTADA INITIO and/or ARISTADA may cause serious side effects including: (continued)

- **Falls.** ARISTADA INITIO and ARISTADA may make you sleepy or dizzy when standing which may make you at risk for falls and related injuries.
- **Low white blood cell count**
- **Seizures (convulsions)**

Please see additional Important Safety Information throughout and full Prescribing Information for **ARISTADA INITIO** and **ARISTADA**, including Boxed Warning and Medication Guides for **ARISTADA INITIO** and **ARISTADA**.

Looking after YOUR LOVED ONE

Living with schizophrenia can be challenging for both you and them. As a caregiver, you do so much to help them. It's a big job.

Talking to your loved one

Here are some things to consider discussing with them:

- Encourage them to feel that they can speak up and advocate for themselves
- Validate their worries and fears
 - Validating sounds like:
“That must be [scary, stressful, etc]. Would it be helpful if...?”
- Make sure you know and understand the side effects of their treatment
- Make sure you and/or your loved one is/are prepared to discuss treatment options with their healthcare provider

The information listed herein is not intended as a substitute for professional medical advice or judgment. Always talk to a healthcare professional.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA *(continued)*

- **Problems controlling your body temperature.**
Avoid becoming too hot or dehydrated. **Do not** exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. **Do not** wear too much clothing or heavy clothing. Drink plenty of water.
- **Difficulty swallowing**

The most common side effects of ARISTADA INITIO and ARISTADA include restlessness or feeling like you need to move (akathisia). These are not all the possible side effects of ARISTADA INITIO and ARISTADA. You should tell your healthcare provider about any side-effects you have.

Do not drive, operate hazardous machinery, or do other dangerous activities until you know how ARISTADA INITIO and ARISTADA affect you. ARISTADA INITIO and ARISTADA may affect your judgment, thinking or motor skills.

Taking care of YOURSELF

It can be a lot to balance work, family, and caregiving. Take some time to look after yourself. Here are some areas that could help:



RECHARGE your batteries

You can't pour from an empty cup! Find time to take part in activities that make you happy—and enjoy them without feeling guilty.



MANAGE your stress

Reduce your levels of stress by getting enough sleep each night, eating well, limiting your intake of alcoholic drinks, and avoiding situations that you know may cause stress for you and the person you care for.



NOTICE the positive

Focus on the day and the joy you can find in it. The little things that go right can add up to a positive experience. If you keep a journal, make sure to write down 1 good thing each day.



Live HEALTHY

Make sure you are taking care of yourself and seeing your healthcare provider for your annual checkups.



Be MINDFUL of your feelings

Being a caregiver can be overwhelming. It's important to notice these feelings without passing judgment and then refocus on the job at hand.



REACH out for support

You are not alone. Don't be afraid to ask for help. Join a support program for caregivers in your area.

Find additional tips on taking care of yourself and your loved one—plus, hear from other caregivers like you—by visiting our website.



aristada.com/caregiver-information

Please see additional Important Safety Information throughout and full Prescribing Information for ARISTADA INITIO and ARISTADA, including Boxed Warning and Medication Guides for ARISTADA INITIO and ARISTADA.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA

ARISTADA INITIO®
aripiprazole lauroxil
extended-release injectable suspension

675 mg

ARISTADA®
aripiprazole lauroxil
extended-release injectable suspension

441 mg 662 mg 882 mg 1064 mg

INDICATION and IMPORTANT SAFETY INFORMATION for ARISTADA INITIO® (aripiprazole lauroxil) and ARISTADA® (aripiprazole lauroxil) extended-release injectable suspension, for intramuscular use

INDICATION

ARISTADA INITIO is a prescription medicine given as a **one-time injection** and is used in combination with oral aripiprazole to start ARISTADA treatment, or re-start ARISTADA treatment after a missed dose, when ARISTADA is used for the treatment of schizophrenia in adults.

ARISTADA is a prescription medicine given by injection by a healthcare professional and used to treat schizophrenia in adults.

It is not known if ARISTADA INITIO and/or ARISTADA are safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION

Elderly people with dementia-related psychosis are at increased risk of death when treated with antipsychotic medicines including ARISTADA INITIO and ARISTADA. ARISTADA INITIO and ARISTADA are not for the treatment of people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia).

Contraindication: Do not receive ARISTADA INITIO or ARISTADA if you are allergic to aripiprazole or any of the ingredients in ARISTADA. Allergic reactions to aripiprazole have ranged from rash, hives and itching to anaphylaxis, which may include difficulty breathing, tightness in the chest, and swelling of the mouth, face, lips, or tongue.

ARISTADA INITIO and/or ARISTADA may cause serious side effects including:

- **Cerebrovascular problems (including stroke) in elderly people with dementia-related psychosis that can lead to death.**

Please see additional Important Safety Information throughout and full Prescribing Information for ARISTADA INITIO and ARISTADA, including Boxed Warning and Medication Guides for ARISTADA INITIO and ARISTADA.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)

- **Neuroleptic malignant syndrome (NMS), a serious condition that can lead to death.** Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS:
 - high fever
 - stiff muscles
 - confusion
 - sweating
 - changes in pulse, heart rate, and blood pressure
- **Uncontrolled body movements (tardive dyskinesia).** ARISTADA INITIO and ARISTADA may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop receiving ARISTADA. Tardive dyskinesia may also start after you stop receiving ARISTADA.
- **Problems with your metabolism such as:**
 - **High blood sugar (hyperglycemia).** Increases in blood sugar can happen in some people who receive ARISTADA INITIO and/or ARISTADA. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes), your healthcare provider should check your blood sugar before you receive ARISTADA INITIO and/or ARISTADA and during your treatment with ARISTADA.
 - **Call your healthcare provider if you have any of these symptoms of high blood sugar:**
 - feel very thirsty
 - need to urinate more than usual
 - feel very hungry
 - feel weak or tired
 - feel sick to your stomach
 - feel confused, or your breath smells fruity
 - **Increased fat levels (cholesterol and triglycerides) in your blood**
 - **Weight gain.** You and your healthcare provider should check your weight regularly.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)

- **Unusual and uncontrollable (compulsive) urges.** Some people taking aripiprazole have had strong unusual urges such as gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having unusual strong urges, talk to your healthcare provider.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** ARISTADA INITIO and ARISTADA may make you sleepy or dizzy when standing which may make you at risk for falls and related injuries.
- **Low white blood cell count**
- **Seizures (convulsions)**
- **Problems controlling your body temperature.** Avoid becoming too hot or dehydrated. **Do not** exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. **Do not** wear too much clothing or heavy clothing. Drink plenty of water.
- **Difficulty swallowing**

The most common side effects of ARISTADA INITIO and ARISTADA include restlessness or feeling like you need to move (akathisia). These are not all the possible side effects of ARISTADA INITIO and ARISTADA. You should tell your healthcare provider about any side-effects you have.

Do not drive, operate hazardous machinery, or do other dangerous activities until you know how ARISTADA INITIO and ARISTADA affect you. ARISTADA INITIO and ARISTADA may affect your judgment, thinking or motor skills.

Before receiving ARISTADA INITIO and ARISTADA tell your healthcare provider about all of your medical conditions, including if you:

- have not taken Abilify®, Abilify Maintena®, or any aripiprazole product before
- have or had heart problems or a stroke
- have diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you receive ARISTADA INITIO and/or ARISTADA and during your treatment with ARISTADA
- have or had low or high blood pressure
- have or had seizures (convulsions)

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA (continued)

- have or had a low white blood cell count
- have problems that may affect you receiving an injection in your buttocks or your arm
- are pregnant or plan to become pregnant. It is not known if ARISTADA INITIO and/or ARISTADA will harm your unborn baby. If you become pregnant while taking ARISTADA INITIO and/or ARISTADA, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388, or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>
- are breastfeeding or plan to breastfeed. ARISTADA INITIO and/or ARISTADA can pass into your breast milk. It is not known if it may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you receive ARISTADA INITIO and/or ARISTADA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ARISTADA INITIO and ARISTADA and other medicines may affect each other causing possible serious side effects. Do not start or stop any medicines after you receive ARISTADA INITIO and ARISTADA without talking to your healthcare provider first.

If you have any questions about your health or medicines, talk to your healthcare provider. You are encouraged to report all side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full Prescribing Information for [ARISTADA INITIO](#) and [ARISTADA](#), including Boxed Warning and Medication Guides for [ARISTADA INITIO](#) and [ARISTADA](#).

The Alkermes logo features the word "Alkermes" in a bold, italicized, black serif font. A red swoosh underline is positioned beneath the letters "A", "l", and "k". A registered trademark symbol (®) is located at the top right of the word.

ARISTADA
INITIO®
aripiprazole lauroxil
extended-release injectable suspension

675 mg

ARISTADA®
aripiprazole lauroxil
extended-release injectable suspension

441 mg

662 mg

882 mg

1064 mg

FOR CAREGIVERS
OF ADULTS WITH
SCHIZOPHRENIA

Consider a treatment
for your loved one that's

1 INJECTION
EVERY 2 MONTHS*

(ARISTADA 1064 mg)

Other dosing options available

For more information and real stories from people
living with schizophrenia, visit [ARISTADA.com](https://www.aristada.com)

*After establishing tolerability to oral aripiprazole and
completing starting regimen.

INDICATION

ARISTADA INITIO® (aripiprazole lauroxil) is a prescription medicine given as a **one-time injection** and is used in combination with oral aripiprazole to start ARISTADA® (aripiprazole lauroxil) treatment, or re-start ARISTADA treatment after a missed dose, when ARISTADA is used for the treatment of schizophrenia in adults.

ARISTADA is a prescription medicine given by injection by a healthcare professional and used to treat schizophrenia in adults.

It is not known if ARISTADA INITIO and/or ARISTADA are safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION FOR ARISTADA INITIO AND ARISTADA

Elderly people with dementia-related psychosis are at increased risk of death when treated with antipsychotic medicines including ARISTADA INITIO and ARISTADA. ARISTADA INITIO and ARISTADA are not for the treatment of people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia).

Please see additional Important Safety Information throughout and full Prescribing Information for [ARISTADA INITIO](#) and [ARISTADA](#), including Boxed Warning and Medication Guides for [ARISTADA INITIO](#) and [ARISTADA](#).

ALKERMES® is a registered trademark of Alkermes, Inc. ARISTADA® and logo, and ARISTADA INITIO®, are registered trademarks of Alkermes Pharma Ireland Limited, used by Alkermes, Inc., under license. ABILIFY® is a trademark of Otsuka Pharmaceutical Company and ABILIFY MAINTENA® is a trademark of Otsuka Pharmaceutical Co., Ltd. ©2022 Alkermes, Inc. All rights reserved. ARI-004645